

Exponential deficiency of convolutions of densities

Iosif Pinelis*

*Department of Mathematical Sciences
Michigan Technological University
Houghton, Michigan 49931, USA
E-mail: ipinelis@mtu.edu*

Abstract: If a probability density $p(\mathbf{x})$ ($\mathbf{x} \in \mathbb{R}^k$) is bounded and $R(t) := \int \mathbf{e}^{\langle \mathbf{x}, t\mathbf{u} \rangle} p(\mathbf{x}) d\mathbf{x} < \infty$ for some linear functional $\mathbf{u}$ and all $t \in (0, 1)$, then, for each $t \in (0, 1)$ and all large enough n , the n -fold convolution of the t -tilted density $\tilde{p}_t(\mathbf{x}) := e^{\langle \mathbf{x}, t\mathbf{u} \rangle} p(\mathbf{x}) / R(t)$ is bounded. This is a corollary of a general, “non-i.i.d.” result, which is also shown to enjoy a certain optimality property. Such results are useful for saddle-point approximations.

AMS 2000 subject classifications: Primary 60E05, 60E10; secondary 60F10, 62E20, 60E15.

Keywords and phrases: probability density, saddle-point approximation, sums of independent random variables/vectors, convolution, exponential integrability, boundedness, tilting, exponential families.

Contents

Contents	1
1 Introduction	2
2 Statements of the results	2
3 Discussion	4
4 Proofs	4
References	11

*Supported by NSF grant DMS-0805946

1. Introduction

Let $\mathbf{X}$ be a random vector in $\mathbb{R}^k$ such that

$$M := \mathbb{E} e^{\lambda \mathbf{e} \mathbf{X}} < \infty \quad (1.1)$$

for some unit vector $\mathbf{e} \in \mathbb{R}^k$ and some $\lambda \in (0, \infty)$; here the juxtaposition $\mathbf{e} \mathbf{x}$ denotes the Euclidean scalar product of vectors $\mathbf{e}$ and $\mathbf{x}$ in $\mathbb{R}^k$. By Chebyshev's inequality, the exponential integrability condition (1.1) implies the tail estimate

$$\mathbb{P}(\mathbf{e} \mathbf{X} \geq x) \leq M e^{-\lambda x} \quad \text{for all } x \in \mathbb{R}. \quad (1.2)$$

Vice versa, for any given $\lambda_0 \in (0, \infty]$ one has the following: if (1.2) holds for each $\lambda \in [0, \lambda_0)$ and some $M = M(\lambda) \in (0, \infty)$, then $\mathbb{E} e^{\lambda \mathbf{e} \mathbf{X}} < \infty$ for each $\lambda \in [0, \lambda_0)$.

Suppose also that (the distribution of) $\mathbf{X}$ has a density p (relative to the Lebesgue measure) such that, for some $\mu \in [0, \lambda)$ and some $C \in (0, \infty)$,

$$p(\mathbf{x}) \leq C e^{-\mu \mathbf{e} \mathbf{x}} \quad \text{for all } \mathbf{x} \in \mathbb{R}^k. \quad (1.3)$$

Note also that, if $\mu = 0$, then condition (1.3) simply means that the density p is bounded.

If p is varying regularly enough in an appropriate sense then, given the condition (1.1), one will have (1.3) for $\mu = \lambda$; that is, one will have an exact “local” counterpart to the “integral” upper bound (1.2). The difference

$$\varepsilon := \lambda - \mu$$

(between the largest possible λ and μ for which (1.1) and (1.3) will still hold) may therefore be referred to as the (exponential) “deficiency” of the density p , which is a measure of its irregularity.

The main result of this paper implies that the deficiency decreases fast under convolution: starting with condition (1.3) for p with $\mu = \lambda - \varepsilon$, one has this condition for the n -fold convolution p^{*n} (in place of p) with $\mu = \lambda - \varepsilon/n$; that is, for the n -fold convolution, the deficiency is n times as small as the original one. More generally, it is proved that, for any probability densities $p_1, \dots, p_n$ on $\mathbb{R}^k$ satisfying the exponential integrability condition with the same λ and with respective deficiencies $\varepsilon_1, \dots, \varepsilon_n$, the deficiency of the convolution $p_1 * \dots * p_n$ is no greater than $\varepsilon^\sharp/n$, where $\varepsilon^\sharp$ stands for the harmonic mean of the original deficiencies $\varepsilon_1, \dots, \varepsilon_n$. Moreover, it is shown that this bound, $\varepsilon^\sharp/n$, cannot be improved.

2. Statements of the results

Let $\mathbf{X}_1, \dots, \mathbf{X}_n$ be any independent random vectors in $\mathbb{R}^k$, with densities $p_1, \dots, p_n$.

Assume the following conditions:

$$M_i := \mathbb{E} e^{\lambda \mathbf{e} \mathbf{X}_i} = \int_{\mathbb{R}^k} e^{\lambda \mathbf{e} \mathbf{x}} p_i(\mathbf{x}) d\mathbf{x} < \infty \quad (2.1)$$

and

$$p_i(\mathbf{x}) \leq C_i e^{-\mu_i \mathbf{e}\mathbf{x}} \quad (2.2)$$

for some C_i 's in $(0, \infty)$, some μ_i 's in $[0, \lambda)$, all $i \in \{1, \dots, n\}$, and all $\mathbf{x} \in \mathbb{R}^k$. Consider the convolution

$$p^{(n)} := p_1 * \dots * p_n, \quad (2.3)$$

which is the density of the sum $\mathbf{X}_1 + \dots + \mathbf{X}_n$.

Theorem 2.1. *There exists a finite constant K_n , which depends only on the numbers n , λ , μ_i , M_i , and C_i , such that*

$$p^{(n)}(\mathbf{x}) \leq K_n e^{-(\lambda - \varepsilon^{(n)}) \mathbf{e}\mathbf{x}} \quad \text{for all } \mathbf{x} \in \mathbb{R}^k, \quad (2.4)$$

where

$$\varepsilon^{(n)} := \frac{1}{\frac{1}{\varepsilon_1} + \dots + \frac{1}{\varepsilon_n}} \quad \text{and} \quad \varepsilon_i := \lambda - \mu_i > 0. \quad (2.5)$$

The necessary proofs will be given in Section 4.

Note that $\varepsilon^{(n)} = \varepsilon^\# / n$, where $\varepsilon^\#$ denotes the harmonic mean of $\varepsilon_1, \dots, \varepsilon_n$. One may also note that $\varepsilon^{(n)} < \min(\varepsilon_1, \dots, \varepsilon_n)$.

It turns out that the coefficient $\lambda - \varepsilon^{(n)}$ in the exponent in the bound (2.4) is the best possible:

Proposition 2.2. *For any natural k and n , any $\lambda \in (0, \infty)$, and any μ_i 's in $[0, \lambda)$, the estimate (2.4) will fail to hold if the number $\varepsilon^{(n)}$ given by (2.5) is replaced by any smaller number.*

From Theorem 2.1, one immediately obtains the particular “i.i.d.” case:

Corollary 2.3. *If conditions (1.1) and (1.3) hold, then for each natural n there exists a constant K_n , which depends only on the numbers n , λ , μ , M , and C , such that*

$$p^{*n}(\mathbf{x}) \leq K_n e^{-(\lambda - \varepsilon/n) \mathbf{e}\mathbf{x}} \quad \text{for all } \mathbf{x} \in \mathbb{R}^k, \quad (2.6)$$

where $\varepsilon := \lambda - \mu$.

It follows from Proposition 2.2 that the coefficient $\lambda - \varepsilon/n$ in the exponent in the bound (2.6) is the best possible.

In turn, Corollary 2.3 yields

Corollary 2.4. *If conditions (1.1) and (1.3) hold, then for each $t \in (0, \lambda)$ there exists a natural number n_t such that for all natural $n \geq n_t$ the n -fold convolution $\tilde{p}_t^{*n}$ of the t -tilted density*

$$\tilde{p}_t(\mathbf{x}) := \frac{e^{t \mathbf{e}\mathbf{x}} p(\mathbf{x})}{\mathbf{E} e^{t \mathbf{e}\mathbf{X}}} \quad (\mathbf{x} \in \mathbb{R}^k) \quad (2.7)$$

is bounded.

In fact, in Corollary 2.4 one may take $n_t = \lceil \frac{\lambda - \mu}{\lambda - t} \rceil$.

Corollary 2.4 can be rewritten as

Corollary 2.5. *If conditions (1.1) and (1.3) hold, then for each $t \in (0, \lambda)$ there exists some $\gamma_t \in (0, \infty)$ such that for all $\gamma \geq \gamma_t$*

$$\int_{\mathbb{R}^k} |\tilde{f}_t(\mathbf{s})|^\gamma d\mathbf{s} < \infty, \quad (2.8)$$

where $\tilde{f}_t(\mathbf{s}) := \int_{\mathbb{R}^k} e^{i\mathbf{s}\mathbf{x}} \tilde{p}_t(\mathbf{x}) d\mathbf{x}$, the characteristic function of the t -tilted density $\tilde{p}_t$; here, of course, i stands for the imaginary unit.

Remark 2.6. In applications, one may of course assume the “grouping”: $\mathbf{X}_j = \mathbf{Y}_{m_{j-1}+1} + \dots + \mathbf{Y}_{m_j}$ for $j = 1, \dots, n$, where $0 = m_0 < m_1 < \dots$ and the $\mathbf{Y}$ ’s are independent random vectors, whose distributions may themselves not have a density. Then the densities $p_1, \dots, p_n$ as in Theorem 2.1 will be the densities of the convolutions of the distributions of the corresponding $\mathbf{Y}$ ’s.

3. Discussion

The condition of the boundedness of the n -fold convolution $\tilde{p}_t^{*n}$ of the tilted density $\tilde{p}_t$ or, equivalently, the condition (2.8) of the absolute integrability of the corresponding “tilted” characteristic function is needed to derive saddle-point approximations. Surveys of literature on such approximations are given e.g. in [3, 9]; for more recent work see e.g. [5, 10].

In the context of saddle-point approximations, the tilting is sometimes described as imbedding the original density p into the exponential family (2.7). The condition of the boundedness of $\tilde{p}_t^{*m}$ for *all* relevant values of the tilting parameter t and all large enough m appears to be usually imposed outright; see e.g. Barndorff-Nielsen and Cox [1, page 298, condition c]; Lugannani and Rice [7, page 481, condition (ii)] impose an even stronger condition, requiring (for $k = 1$) that $|\tilde{f}_t(s)| = O((1 + |s|)^{-\gamma})$ for some $\gamma > 0$ and all $s \in \mathbb{R}$.

On the other hand, Corollaries 2.4 and 2.5 together with Remark 2.6 show that one need *a priori* require the boundedness of $\tilde{p}_t^{*m}$ only for $t = 0$ and some natural m , that is, only for some convolution p^{*m} of the original, un-tilted density p ; then $\tilde{p}_t^{*m}$ will necessarily be bounded for all t in the interval $[0, \lambda)$ and all large enough m .

The considerations presented above in this section constituted the original motivation for the present work. The proof of Proposition 2.2 (given in the next section) shows that probability densities with the deficiencies most resistant to convolution are mixtures of infinitely many mutually (almost) singular densities, spaced regularly enough (see Fig. 1 on page 7). Such “exponentially deficient” distributions can be contrasted with the well-studied classes of regularly behaving distributions with nearly exponential tails; see e.g. [4, 6, 8].

4. Proofs

Proof of Theorem 2.1. To begin, note that for $n = 1$ the inequality (2.4) with $K_1 := C_1$ is the same as (2.2). Next, a trivial remark is that (2.1) implies

$\mathbb{E} e^{\lambda \mathbf{e}(\mathbf{X}_1 + \dots + \mathbf{X}_{n-1})} = M_1 \cdots M_{n-1} < \infty$. Note also that (2.5) can be rewritten in an additive form, as

$$\frac{1}{\varepsilon^{(n)}} = \frac{1}{\varepsilon_1} + \dots + \frac{1}{\varepsilon_n}.$$

So, by induction, it suffices to prove Theorem 2.1 for $n = 2$. For such a case, let us simplify the notation by writing p and q instead of p_1 and p_2 , M and N instead of M_1 and M_2 , C and D instead of C_1 and C_2 , μ and ν instead of μ_1 and μ_2 , and ε and δ instead of ε_1 and ε_2 .

Next, without loss of generality, $\mathbf{e} = (1, 0, \dots, 0) \in \mathbb{R}^k$. Then, identifying any vector $\mathbf{x} \in \mathbb{R}^k$ with the corresponding pair $(x, \mathbf{y}) \in \mathbb{R} \times \mathbb{R}^{k-1}$, one has $\mathbf{e}\mathbf{x} = x$, so that (2.2) can in this case be rewritten as

$$p(x, \mathbf{y}) \leq C e^{-\mu x} \quad \text{and} \quad q(x, \mathbf{y}) \leq D e^{-\nu x} \quad (4.1)$$

for all $(x, \mathbf{y}) \in \mathbb{R} \times \mathbb{R}^{k-1}$. Also, conditions (2.1) imply

$$\int_x^\infty du \tilde{p}(u) \leq M e^{-\lambda x} \quad \text{and} \quad \int_x^\infty du \tilde{q}(u) \leq N e^{-\lambda x} \quad (4.2)$$

for all $x \in \mathbb{R}$, where

$$\tilde{p}(u) := \int_{\mathbb{R}^k} d\mathbf{v} p(u, \mathbf{v}) \quad \text{and} \quad \tilde{q}(u) := \int_{\mathbb{R}^k} d\mathbf{v} q(u, \mathbf{v})$$

for all $u \in \mathbb{R}$, the densities of the random variables $\mathbf{e}\mathbf{X}_1$ and $\mathbf{e}\mathbf{X}_2$, respectively.

Fix now any $(x, \mathbf{y}) \in \mathbb{R} \times \mathbb{R}^{k-1}$. Take, for a moment, any $\alpha \in (0, 1)$ and let $\beta := 1 - \alpha$. Then

$$(p * q)(x, \mathbf{y}) = \int_{\mathbb{R}} du \int_{\mathbb{R}^{k-1}} d\mathbf{v} p(x - u, \mathbf{y} - \mathbf{v}) q(u, \mathbf{v}) \leq D I_1 + C I_2 \quad (4.3)$$

by (4.1), where

$$\begin{aligned} I_1 &:= \int_{-\infty}^{\alpha x} du \int_{\mathbb{R}^{k-1}} d\mathbf{v} p(x - u, \mathbf{y} - \mathbf{v}) e^{-\nu u} = \int_{-\infty}^{\alpha x} du \tilde{p}(x - u) e^{-\nu u}, \\ I_2 &:= \int_{\alpha x}^\infty du \int_{\mathbb{R}^{k-1}} d\mathbf{v} p(x - u, \mathbf{y} - \mathbf{v}) q(u, \mathbf{v}) = \int_{-\infty}^{\beta x} dz \tilde{q}(x - z) e^{-\mu z}. \end{aligned}$$

Next, in view of (4.2),

$$\begin{aligned} I_1 &= \int_{-\infty}^{\alpha x} du \tilde{p}(x - u) \int_u^\infty \nu dz e^{-\nu z} \\ &= \int_{-\infty}^{\alpha x} \nu dz e^{-\nu z} \int_{-\infty}^z du \tilde{p}(x - u) + \int_{\alpha x}^\infty \nu dz e^{-\nu z} \int_{-\infty}^{\alpha x} du \tilde{p}(x - u) \\ &= \int_{-\infty}^{\alpha x} \nu dz e^{-\nu z} \int_{x-z}^\infty dw \tilde{p}(w) + \int_{\alpha x}^\infty \nu dz e^{-\nu z} \int_{(1-\alpha)x}^\infty dw \tilde{p}(w) \\ &\leq \int_{-\infty}^{\alpha x} \nu dz e^{-\nu z} M e^{-\lambda(x-z)} + \int_{\alpha x}^\infty \nu dz e^{-\nu z} M e^{-\lambda(1-\alpha)x} \\ &= M \frac{\lambda}{\lambda - \nu} e^{-(\lambda - (\lambda - \nu)\alpha)x}. \end{aligned} \quad (4.4)$$

Note that this derivation of the upper bound (4.5) on I_1 is valid only for $\nu \neq 0$. However, if $\nu = 0$, then

$$I_1 = \int_{-\infty}^{\alpha x} du \tilde{p}(x-u) = \int_{(1-\alpha)x}^{\infty} dz \tilde{p}(z) \leq M e^{-\lambda(1-\alpha)x},$$

so that the bound (4.5) on I_1 holds for $\nu = 0$ as well. Recall now that $\varepsilon = \lambda - \mu$ and $\delta = \lambda - \nu$, and choose $\alpha := \frac{\varepsilon}{\varepsilon + \delta}$. Then (4.5) can be rewritten as

$$I_1 \leq M \frac{\lambda}{\delta} e^{-(\lambda - \varepsilon^{(2)})x}, \quad (4.6)$$

with $\varepsilon^{(2)} = \frac{\varepsilon\delta}{\varepsilon + \delta} = \frac{1}{1/\varepsilon + 1/\delta}$, in accordance with the definition (2.5) of $\varepsilon^{(n)}$. Quite similarly,

$$I_2 \leq N \frac{\lambda}{\varepsilon} e^{-(\lambda - \varepsilon^{(2)})x}. \quad (4.7)$$

Collecting now (4.3), (4.6), and (4.7), one sees that

$$(p * q)(x, \mathbf{y}) \leq K_2 e^{-(\lambda - \varepsilon^{(2)})x} \quad (4.8)$$

for some constant K_2 depending only on λ, μ, M, N, C , and D , and for all $(x, \mathbf{y}) \in \mathbb{R} \times \mathbb{R}^{k-1}$. Thus, Theorem 2.1 is proved for $n = 2$ and, thereby, for all natural n . $\square$

The proof of Proposition 2.2 rests on Lemma 4.1 below. To state the lemma, for any $\lambda \in (0, \infty)$ and $\varepsilon \in (0, \lambda]$ introduce the class $\mathcal{P}_{\lambda, \varepsilon}$ of all probability densities p on $\mathbb{R}$ such that

- (i) $\int_{\mathbb{R}} e^{\lambda x} p(x) dx < \infty$ and
- (ii) $p(x) \geq c p_{\lambda, \varepsilon, \kappa, \alpha}(x)$ for some $c \in (0, \infty)$, $\kappa \in (0, \infty)$, $\alpha \in (\frac{1}{2}, \infty)$, and all $x \in \mathbb{R}$, where

$$p_{\lambda, \varepsilon, \kappa, \alpha}(x) := \sum_{j=-\infty}^{\infty} W_j(x), \quad (4.9)$$

$$W_j(x) := W_{j; \lambda, \varepsilon, \kappa, \alpha}(x) := w_j f_{j, \kappa e^{-\varepsilon|j|}}(x), \quad (4.10)$$

$$w_j := w_{j; \lambda, \alpha} := \frac{e^{-\lambda|j|}}{(j^2 + 1)^\alpha}, \quad (4.11)$$

$$f_{a, b}(x) := \frac{1}{b} \varphi\left(\frac{x-a}{b}\right), \quad \varphi(u) := \frac{1}{\sqrt{2\pi}} e^{-u^2/2};$$

of course, $f_{a, b}$ is the density of the normal distribution with mean a and variance b^2 .

(One could similarly, and even a little more easily, deal with the “asymmetric” version of the class $\mathcal{P}_{\lambda, \varepsilon}$, having $\sum_{j=-\infty}^{\infty}$ in (4.9) replaced by $\sum_{j=0}^{\infty}$.)

Lemma 4.1. *Take any $\lambda \in (0, \infty)$, $\varepsilon \in (0, \lambda]$, $\kappa \in (0, \infty)$, and $\alpha \in (\frac{1}{2}, \infty)$.*

(I) There exists some $c_{\lambda,\varepsilon,\kappa,\alpha} \in (0, \infty)$ such that $\tilde{p}_{\lambda,\varepsilon,\kappa,\alpha} := \frac{p_{\lambda,\varepsilon,\kappa,\alpha}}{c_{\lambda,\varepsilon,\kappa,\alpha}} \in \mathcal{P}_{\lambda,\varepsilon}$.

In particular, it follows that $\mathcal{P}_{\lambda,\varepsilon} \neq \emptyset$.

(II) There exists some $C = C_{\lambda,\varepsilon,\kappa,\alpha} \in (0, \infty)$ such that for $p = p_{\lambda,\varepsilon,\kappa,\alpha}$ and $\mu := \lambda - \varepsilon$

$$p(x) \leq C e^{-\mu x} \quad \text{for all } x \in \mathbb{R}. \quad (4.12)$$

(III) For any $p \in \mathcal{P}_{\lambda,\varepsilon}$ and any $C \in (0, \infty)$, relation (4.12) does not hold with any $\mu^\diamond \in (\lambda - \varepsilon, \infty)$ in place of μ .

(IV) In addition to ε , take any $\delta \in (0, \lambda]$. Then, for any $p \in \mathcal{P}_{\lambda,\varepsilon}$ and $q \in \mathcal{P}_{\lambda,\delta}$, one has $p * q \in \mathcal{P}_{\lambda,\tilde{\varepsilon}}$, where

$$\tilde{\varepsilon} := \frac{1}{\frac{1}{\varepsilon} + \frac{1}{\delta}} = \frac{\varepsilon\delta}{\varepsilon + \delta}. \quad (4.13)$$

The (symmetric about 0) probability density $\tilde{p}_{\lambda,\varepsilon,\kappa,\alpha}$ as in part (I) of this lemma is illustrated here:




FIG 1. Graphs $\{(x, p(x)) : 0 < x < 7.5\}$ and $\{(x, \ln p(x)) : 0 < x < 7.5\}$ for $p = \tilde{p}_{\lambda,\varepsilon,\kappa,\alpha}$ with $\lambda = 0.55$, $\varepsilon = 0.50$, $\kappa = 0.9$, and $\alpha = 0.6$

Let us postpone the proof of Lemma 4.1, which is somewhat long, and proceed now to the proof of Proposition 2.2.

Proof of Proposition 2.2. Take indeed any natural k and n , any $\lambda \in (0, \infty)$, and any $\mu_1, \dots, \mu_n$ in $[0, \lambda)$. In accordance with (2.5), let $\varepsilon_i := \lambda - \mu_i$, so that $\varepsilon_i \in (0, \lambda]$ for all $i = 1, \dots, n$. For each $i = 1, \dots, n$, take any density $q_i \in \mathcal{P}_{\lambda,\varepsilon_i}$ such that

$$q_i(x) \leq C_i e^{-\mu_i x} \quad (4.14)$$

for some finite positive real constant C_i and all $x \in \mathbb{R}$; by parts (I) and (II) of Lemma 4.1, such q_i 's do exist.

As in the proof of Theorem 2.1, let $\mathbf{e} = (1, 0, \dots, 0) \in \mathbb{R}^k$ and identify any vector $\mathbf{x} \in \mathbb{R}^k$ with $(x, \mathbf{y}) \in \mathbb{R} \times \mathbb{R}^{k-1}$. Then, for each $i = 1, \dots, n$, introduce the densities

$$p_i(\mathbf{x}) = p_i(x, \mathbf{y}) := q_i(x) \varphi_{k-1}(\mathbf{y}) \quad (4.15)$$

for all $\mathbf{x} = (x, \mathbf{y}) \in \mathbb{R} \times \mathbb{R}^{k-1}$, where $\varphi_{k-1}(\mathbf{y}) := (2\pi)^{-(k-1)/2} e^{-\mathbf{y}\mathbf{y}/2}$ for all $\mathbf{y} \in \mathbb{R}^{k-1}$; then

$$\int_{\mathbb{R}^k} e^{\lambda \mathbf{e}\mathbf{x}} p_i(\mathbf{x}) d\mathbf{x} = \int_{\mathbb{R}} e^{\lambda x} q_i(x) dx < \infty,$$

since $q_i \in \mathcal{P}_{\lambda, \varepsilon_i}$; also, by (4.14),

$$p_i(\mathbf{x}) = q_i(x) \varphi_{k-1}(\mathbf{y}) \leq (2\pi)^{-(k-1)/2} q_i(x) \leq C_i e^{-\mu_i x}$$

for all $\mathbf{x} = (x, \mathbf{y}) \in \mathbb{R} \times \mathbb{R}^{k-1}$. So, conditions (2.1) and (2.2) hold.

Next, introduce

$$q^{(n)} := q_1 * \cdots * q_n,$$

so that, by (2.3) and (4.15),

$$p^{(n)}(\mathbf{x}) = p^{(n)}(x, \mathbf{y}) = q^{(n)}(x) \varphi_{k-1}^{*n}(\mathbf{y}) \quad (4.16)$$

for all $\mathbf{x} = (x, \mathbf{y}) \in \mathbb{R} \times \mathbb{R}^{k-1}$. Moreover, recalling the conditions $q_i \in \mathcal{P}_{\lambda, \varepsilon_i}$ for $i = 1, \dots, n$ and using part (IV) of Lemma 4.1, by induction one concludes that $q^{(n)} \in \mathcal{P}_{\lambda, \varepsilon^{(n)}}$.

Now, to obtain a contradiction, assume that (2.4) holds with some “deficiency” $\varepsilon^\diamond$ in place of $\varepsilon^{(n)}$ such that $\varepsilon^\diamond < \varepsilon^{(n)}$. Then, by (4.16), for $\mu^\diamond := \lambda - \varepsilon^\diamond$

$$q^{(n)}(x) \varphi_{k-1}^{*n}(\mathbf{0}) \leq K_n e^{-\mu^\diamond x}$$

for some constant K_n and all $x \in \mathbb{R}$. But this contradicts part (III) of Lemma 4.1, since $\mu^\diamond \in (\lambda - \varepsilon^{(n)}, \infty)$, $q^{(n)} \in \mathcal{P}_{\lambda, \varepsilon^{(n)}}$, and $\varphi_{k-1}^{*n}(\mathbf{0}) = (2\pi n)^{-(k-1)/2} > 0$. This concludes the proof of Proposition 2.2, except that one still needs to prove Lemma 4.1. $\square$

Proof of Lemma 4.1.

(I) Obviously, $p_{\lambda, \varepsilon, \kappa, \alpha} > 0$ and $c_{\lambda, \varepsilon, \kappa, \alpha} := \int_{\mathbb{R}} p_{\lambda, \varepsilon, \kappa, \alpha}(x) dx = \sum_{j=-\infty}^{\infty} w_j < \infty$ for any $\lambda \in (0, \infty)$, $\varepsilon \in (0, \lambda]$, $\kappa \in (0, \infty)$, and $\alpha \in (\frac{1}{2}, \infty)$. So, $\tilde{p}_{\lambda, \varepsilon, \kappa, \alpha}$ is a probability density. Moreover,

$$\int_{\mathbb{R}} e^{\lambda x} p_{\lambda, \varepsilon, \kappa, \alpha}(x) dx = \sum_{j=-\infty}^{\infty} w_j \exp\left(\lambda j + \frac{1}{2} \lambda^2 \kappa^2 e^{-2\varepsilon|j|}\right) \leq \sum_{j=-\infty}^{\infty} \frac{e^{\lambda^2 \kappa^2 / 2}}{(j^2 + 1)^\alpha} < \infty. \quad (4.17)$$

Thus, part (I) of Lemma 4.1 is verified.

(II) Note that

$$W_j(x) = \frac{1}{\kappa \sqrt{2\pi}} \frac{e^{-(\lambda - \varepsilon)|j|}}{(j^2 + 1)^\alpha} \exp\left(-\frac{(x - j)^2 e^{2\varepsilon|j|}}{2\kappa^2}\right). \quad (4.18)$$

Hence and because $\varepsilon \in (0, \lambda]$, one has $p_{\lambda, \varepsilon, \kappa, \alpha}(x) \leq C := \sum_{j=-\infty}^{\infty} W_j(j) < \infty$ for all $x \in \mathbb{R}$. So, $p_{\lambda, \varepsilon, \kappa, \alpha}(x) \leq C e^{-\mu x}$ for all $x \in (-\infty, 0]$; that is, (4.12) holds for $p = p_{\lambda, \varepsilon, \kappa, \alpha}$, $\mu = \lambda - \varepsilon$, and all $x \in (-\infty, 0]$.

Take now any $x \in (0, \infty)$. Introduce $j_x := \lfloor x \rfloor$, so that $0 \leq j_x \leq x < j_x + 1$ and for $j \geq j_x$ one has $|j| = j > x - 1$. Then, in view of (4.18),

$$\sum_{j=j_x}^{\infty} W_j(x) \leq \sum_{j=j_x}^{\infty} W_j(j) \leq \frac{e^{-(\lambda-\varepsilon)(x-1)}}{\kappa\sqrt{2\pi}} \sum_{j=0}^{\infty} \frac{1}{(j^2+1)^\alpha} = c_1 e^{-(\lambda-\varepsilon)x}; \quad (4.19)$$

in this proof of part (II) of the lemma, let $c_1, c_2, \dots$ denote finite positive constants depending only on $\lambda, \varepsilon, \kappa, \alpha$. Next, for $r_x := \lceil \kappa\sqrt{2(\lambda-\varepsilon)x} \rceil$ and $j \in (-\infty, j_x - r_x]$, one has $x - j \geq r_x$, whence

$$\sum_{j=-\infty}^{j_x - r_x} W_j(x) \leq \sum_{j=-\infty}^{j_x - r_x} W_j(j) \exp - \frac{r_x^2}{2\kappa^2} \leq c_2 \exp - \frac{r_x^2}{2\kappa^2} \leq c_2 e^{-(\lambda-\varepsilon)x}. \quad (4.20)$$

Further, for $j \in [j_x - r_x + 1, j_x - 1]$ one has $x - j \geq 1$ and $|j| \geq j \geq j_x - r_x + 1 > x - r_x \geq \frac{x}{2} - c_3$, whence

$$\begin{aligned} \sum_{j=j_x - r_x + 1}^{j_x - 1} W_j(x) &\leq \sum_{j=j_x - r_x + 1}^{j_x - 1} W_j(j) \exp - \frac{e^{2\varepsilon(x-r_x)}}{2\kappa^2} \leq c_4 \exp - \frac{e^{\varepsilon(x-2c_3)}}{2\kappa^2} \\ &\leq c_5 e^{-(\lambda-\varepsilon)x}. \end{aligned} \quad (4.21)$$

So, by (4.9), (4.19), (4.20), and (4.21), the relation (4.12) (with $\mu = \lambda - \varepsilon$) holds for $p = p_{\lambda, \varepsilon, \kappa, \alpha}$ and all $x \in (0, \infty)$ as well. This completes the verification of part (II) of the lemma.

(III) Take any $p \in \mathcal{P}_{\lambda, \varepsilon}$, so that $p \geq c p_{\lambda, \varepsilon, \kappa, \alpha}$ for some $c \in (0, \infty)$, $\kappa \in (0, \infty)$, and $\alpha \in (\frac{1}{2}, \infty)$. Then

$$p(j) \geq c p_{\lambda, \varepsilon, \kappa, \alpha}(j) \geq c W_j(j) = \frac{c}{\kappa\sqrt{2\pi}} \frac{e^{-(\lambda-\varepsilon)j}}{(j^2+1)^\alpha} > C e^{-\mu^\diamond j}$$

for any $\mu^\diamond \in (\lambda - \varepsilon, \infty)$, any $C \in (0, \infty)$, and all large enough natural j . This proves part (III) of the lemma.

(IV) Take any $p \in \mathcal{P}_{\lambda, \varepsilon}$ and $q \in \mathcal{P}_{\lambda, \delta}$, so that $p \geq c p_{\lambda, \varepsilon, \kappa, \alpha}$ and $q \geq \tilde{c} p_{\lambda, \delta, \xi, \beta}$ for some $c \in (0, \infty)$, $\kappa \in (0, \infty)$, $\alpha \in (\frac{1}{2}, \infty)$, $\tilde{c} \in (0, \infty)$, $\xi \in (0, \infty)$, and $\beta \in (\frac{1}{2}, \infty)$.

Choose for a moment any $m \in \{0, 1, \dots\}$ and let

$$i_m := \left\lfloor m \frac{\delta}{\varepsilon + \delta} \right\rfloor \quad \text{and} \quad j_m := \left\lceil m \frac{\varepsilon}{\varepsilon + \delta} \right\rceil = m - i_m, \quad (4.22)$$

so that $m \frac{\delta}{\varepsilon + \delta} - 1 \leq i_m \leq m \frac{\delta}{\varepsilon + \delta}$ and $m \frac{\varepsilon}{\varepsilon + \delta} \leq j_m \leq m \frac{\varepsilon}{\varepsilon + \delta} + 1$. Next, introduce

$$\begin{aligned} \sigma_m &:= \sqrt{\kappa^2 e^{-2\varepsilon|i_m|} + \xi^2 e^{-2\delta|j_m|}} = \sqrt{\kappa^2 e^{-2\varepsilon i_m} + \xi^2 e^{-2\delta j_m}}, \\ \zeta &:= \sqrt{\kappa^2 + \xi^2 e^{-2\delta}}, \quad \tilde{\zeta} := \sqrt{\kappa^2 e^{2\varepsilon} + \xi^2}, \end{aligned}$$

and observe that

$$\begin{aligned} \frac{\tilde{\zeta}}{\zeta} &\leq e^{\varepsilon \vee \delta}, \\ \sigma_m^2 &\geq \kappa^2 \exp \left\{ -2\varepsilon m \frac{\delta}{\varepsilon + \delta} \right\} + \xi^2 \exp \left\{ -2\delta \left(m \frac{\varepsilon}{\varepsilon + \delta} + 1 \right) \right\} = \zeta^2 e^{-2\tilde{\varepsilon}m}, \\ \sigma_m^2 &\leq \kappa^2 \exp \left\{ -2\varepsilon \left(m \frac{\delta}{\varepsilon + \delta} - 1 \right) \right\} + \xi^2 \exp \left\{ -2\delta m \frac{\varepsilon}{\varepsilon + \delta} \right\} = \tilde{\zeta}^2 e^{-2\tilde{\varepsilon}m}, \end{aligned}$$

where $\tilde{\varepsilon}$ is as in (4.13). Also, recall that here $m \geq 0$, $i_m \geq 0$, and $j_m \geq 0$. It follows that for all $x \in \mathbb{R}$

$$\begin{aligned} (f_{i_m, \kappa e^{-\varepsilon|i_m|}} * f_{j_m, \xi e^{-\delta|j_m|}})(x) &= f_{m, \sigma_m^2}(x) = \frac{1}{\sigma_m \sqrt{2\pi}} \exp -\frac{(x-m)^2}{2\sigma_m^2} \\ &\geq \frac{\zeta e^{-\tilde{\varepsilon}m}}{\sigma_m} f_{m, \zeta e^{-\tilde{\varepsilon}m}}(x) \geq \frac{\zeta}{\tilde{\zeta}} f_{m, \zeta e^{-\tilde{\varepsilon}m}}(x) \geq e^{-(\varepsilon \vee \delta)} f_{m, \zeta e^{-\tilde{\varepsilon}|m|}}(x). \end{aligned}$$

Quite similarly (or by symmetry), one has

$$f_{i_m, \kappa e^{-\varepsilon|i_m|}} * f_{j_m, \xi e^{-\delta|j_m|}} \geq e^{-(\varepsilon \vee \delta)} f_{m, \zeta e^{-\tilde{\varepsilon}|m|}}$$

for any $m \in \{-1, -2, \dots\}$, letting now $i_m := -i_{-m} = \lceil m \frac{\delta}{\varepsilon + \delta} \rceil$ and $j_m := -j_{-m} = \lfloor m \frac{\varepsilon}{\varepsilon + \delta} \rfloor$, so that still $i_m + j_m = m$.

On recalling the conditions $p \geq c p_{\lambda, \varepsilon, \kappa, \alpha}$, $q \geq \tilde{c} p_{\lambda, \delta, \xi, \beta}$, (4.9)–(4.11), and (4.22), it follows that

$$\begin{aligned} p * q &\geq c \tilde{c} \sum_{m=-\infty}^{\infty} w_{i_m; \lambda, \alpha} w_{j_m; \lambda, \beta} f_{i_m, \kappa e^{-\varepsilon|i_m|}} * f_{j_m, \xi e^{-\delta|j_m|}} \\ &\geq c_1 \sum_{m=-\infty}^{\infty} w_{i_m; \lambda, \alpha} w_{j_m; \lambda, \beta} f_{m, \zeta e^{-\tilde{\varepsilon}|m|}} \\ &\geq c_2 \sum_{m=-\infty}^{\infty} w_{m; \lambda, \alpha + \beta} f_{m, \zeta e^{-\tilde{\varepsilon}|m|}} = c_2 p_{\lambda, \tilde{\varepsilon}, \zeta, \alpha + \beta}, \end{aligned}$$

where c_1 and c_2 are strictly positive constants depending only on $\lambda, \varepsilon, \delta, \kappa, \xi, \alpha, \beta$.

Also, $\int_{\mathbb{R}} e^{\lambda x} (p * q)(x) dx = \int_{\mathbb{R}} e^{\lambda x} p(x) dx \int_{\mathbb{R}} e^{\lambda x} q(x) dx < \infty$. Thus, it has been shown that $p * q \in \mathcal{P}_{\lambda, \tilde{\varepsilon}}$. This completes the verification of part (IV). The lemma is now completely proved. $\square$

Proof of Corollary 2.4. This follows because $\tilde{p}_t^{*n}(\mathbf{x}) = \frac{e^{t \mathbf{ex}} p^{*n}(\mathbf{x})}{(\mathbb{E} e^{t \mathbf{eX}})^n}$ for all $\mathbf{x} \in \mathbb{R}^k$. $\square$

Proof of Corollary 2.5. Take any $t \in (0, \lambda)$. Then, by Corollary 2.4, $\tilde{p}_t^{*n_t}$ is bounded by some constant $K < \infty$. Then, by the Plancherel isometry (see e.g. [2, Theorem 4.2]), for all $\gamma \geq 2n_t$

$$\int_{\mathbb{R}^k} |\tilde{f}_t(\mathbf{s})|^\gamma d\mathbf{s} \leq \int_{\mathbb{R}^k} |\tilde{f}_t(\mathbf{s})|^{2n_t} d\mathbf{s} = (2\pi)^k \int_{\mathbb{R}^k} \tilde{p}_t^{*n_t}(\mathbf{x})^2 d\mathbf{x} \leq (2\pi)^k K < \infty.$$

Vice versa, assume that (2.8) holds for all $\gamma \geq \gamma_t$; then $\tilde{p}_t^{*n}$ is bounded for all natural $n \geq \gamma_t$ by the Fourier inversion formula (see e.g. [2, Theorem 4.1(iv)]), since the characteristic function of $\tilde{p}_t^{*n}$ is $\tilde{f}_t(\mathbf{s})^n$. $\square$

Remark 4.2. Weaker results than the one given by Theorem 2.1 or even Corollary 2.3 (but which still be enough to deduce Corollaries 2.4 and 2.5) can be obtained more simply modulo the Plancherel isometry. Indeed, if conditions (1.1) and (1.3) hold, then

$$\int_{\mathbb{R}^k} |\tilde{f}_t(\mathbf{s})|^2 d\mathbf{s} = (2\pi)^k \int_{\mathbb{R}^k} \tilde{p}_t(\mathbf{x})^2 d\mathbf{x} = \frac{(2\pi)^k}{(\mathbb{E} e^{t \mathbf{e} \mathbf{X}})^2} \int_{\mathbb{R}^k} e^{2t \mathbf{e} \mathbf{x}} p(\mathbf{x})^2 d\mathbf{x} < \infty$$

for $t = \lambda - \varepsilon/2$ and $\varepsilon := \lambda - \mu$, since $e^{2(\lambda - \varepsilon/2) \mathbf{e} \mathbf{x}} p(\mathbf{x})^2 \leq C e^{\lambda \mathbf{e} \mathbf{x}} p(\mathbf{x})$ for all $\mathbf{x} \in \mathbb{R}$. Also, by the Fourier inversion formula, again with $t = \lambda - \varepsilon/2$,

$$\tilde{p}_t^{*2}(\mathbf{x}) \leq (2\pi)^{-k} \int_{\mathbb{R}^k} |\tilde{f}_t(\mathbf{s})|^2 d\mathbf{s} < \infty \quad \text{for all } \mathbf{x} \in \mathbb{R}^k,$$

which yields (2.6) for $n = 2$. Thus, by induction, one can obtain (2.6) for $n = 2^j$, where j is any natural number.

However, it is unclear whether such an approach, via the Plancherel isometry, could be extended to yield Theorem 2.1 or, at least, Corollary 2.3 for all natural n . Anyway, it might be not worthwhile to exert efforts in such a direction, as the direct probabilistic proof of Theorem 2.1 given above is rather simple already and yet produces the best possible bound on the exponential deficiency.

References

- [1] BARNDORFF-NIELSEN, O., AND COX, D. R. Edgeworth and saddle-point approximations with statistical applications. *J. Roy. Statist. Soc. Ser. B* 41, 3 (1979), 279–312. With discussion.
- [2] BHATTACHARYA, R. N., AND RANGA RAO, R. *Normal approximation and asymptotic expansions*. Robert E. Krieger Publishing Co. Inc., Melbourne, FL, 1986. Reprint of the 1976 original.
- [3] DANIELS, H. E. Tail probability approximations. *Internat. Statist. Rev.* 55, 1 (1987), 37–48.
- [4] EMBRECHTS, P., AND GOLDIE, C. M. On convolution tails. *Stochastic Process. Appl.* 13, 3 (1982), 263–278.
- [5] JING, B.-Y., SHAO, Q.-M., AND ZHOU, W. Saddlepoint approximation for Student's t -statistic with no moment conditions. *Ann. Statist.* 32, 6 (2004), 2679–2711.
- [6] KLÜPPELBERG, C. Subexponential distributions and characterizations of related classes. *Probab. Theory Related Fields* 82, 2 (1989), 259–269.
- [7] LUGANNANI, R., AND RICE, S. Saddle point approximation for the distribution of the sum of independent random variables. *Adv. in Appl. Probab.* 12, 2 (1980), 475–490.

- [8] PINELIS, I. F. Asymptotic equivalence of the probabilities of large deviations for sums and maximum of independent random variables. In *Limit theorems of probability theory*, vol. 5 of *Trudy Inst. Mat.* “Nauka” Sibirsk. Otdel., Novosibirsk, 1985, pp. 144–173, 176.
- [9] REID, N. Saddlepoint methods and statistical inference. *Statist. Sci.* 3, 2 (1988), 213–238. With comments and a rejoinder by the author.
- [10] SHAO, Q.-M. Recent progress on self-normalized limit theorems. In *Probability, finance and insurance*. World Sci. Publ., River Edge, NJ, 2004, pp. 50–68.